





Web: www.anbobio.com
www.dijibio.com
Order: order@dijibio.com
Tel: (86) 519 8805 0026

DARPP-32 (Phospho-Thr75) Antibody [#A0007]

- Catalog Number:** A0007
Amount: 50µg/50µl, 100µg/100µl, 200µg/200µl
Swiss-Prot No. : Q9UD71
All Names: Dopamine- and cAMP-regulated neuronal phosphoprotein, IPPD, Neuronal phosphoprotein DARPP-32, PPP1R1B
All Sites: Human:Thr75; Mouse: Thr75; Rat: Thr75
Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage/Stability: Store at -20°C/1 year
Immunogen: The antiserum was produced against synthesized phosphopeptide derived from human DARPP-32 around the phosphorylation site of threonine 75 (A-Y-T^P-P-P).
Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatography using non-phosphopeptide corresponding to the phosphorylation site.
Specificity/Sensitivity: DARPP-32 (phospho-Thr75) antibody detects endogenous levels of DARPP-32 only when phosphorylated at threonine 75.
Reactivity: Human, Mouse, Rat
Applications: WB: 1:500~3000 IHC: 1:50~1:100
ELISA: 1:40000
References: JA Girault, J. Biol. Chem., Dec 1989; 264: 21748 - 21759.
GL Snyder, J. Neurosci., Aug 1992; 12: 3071.
Akinori Nishi, J. Neurosci., Nov 1997; 17: 8147.
Artur Mayerhofer, J. Clin. Endocrinol. Metab., Dec 2000; 85: 4750 - 4757.



Immunohistochemical analysis of paraffin-embedded human brain tissue, using DARPP-32 (phospho-Thr75) antibody.



Western blot analysis of extracts from COS7 cells treated with Forskolin (40nM, 30mins), using DARPP-32 (phospho-Thr75) antibody.

For Research Use Only